

Modified Growth Diagrams, Permutation Pivots, and the BWX Map ϕ^*

Jonathan Bloom
Dartmouth College

Dan Saracino
Colgate University

Abstract

In their paper on Wilf-equivalence for singleton classes, Backelin, West, and Xin introduced a transformation ϕ^* , defined by an iterative process and operating on (all) full rook placements on Ferrers boards. Bousquet-Mélou and Steingrímsson proved the analogue of the main result of Backelin, West, and Xin in the context of involutions, and in so doing they needed to prove that ϕ^* commutes with the operation of taking inverses. The proof of this commutation result was long and difficult, and Bousquet-Mélou and Steingrímsson asked if ϕ^* might be reformulated in such a way as to make this result obvious. In the present paper we provide such a reformulation of ϕ^* , by modifying the growth diagram algorithm of Fomin. This also answers a question of Krattenthaler, who noted that a bijection defined by the unmodified Fomin algorithm obviously commutes with inverses, and asked what the connection is between this bijection and ϕ^* .

1. Introduction

For any permutation $\tau = \tau_1 \tau_2 \dots \tau_r$, let $S_n(\tau)$ denote the set of permutations in S_n that avoid τ , in the sense that they have no subsequence order-isomorphic to τ .

In their paper *Wilf-equivalence for singleton classes* [1], Backelin, West, and Xin prove an important general result about permutations avoiding a single pattern: If $k, \ell \geq 1$ and ρ is a permutation of $\{k+1, \dots, k+\ell\}$, then for every $n \geq k+\ell$ we have $|S_n(12\dots k\rho)| = |S_n(k\dots 1\rho)|$. The key tool in the proof is a map ϕ^* , which operates on a permutation σ as follows: Order the $k\dots 1$ -patterns $\sigma_{i_1} \dots \sigma_{i_k}$ in σ lexicographically (according to the σ_j 's, not the j 's) and let $\phi(\sigma)$ be obtained from σ by taking the smallest $\sigma_{i_1} \dots \sigma_{i_k}$ and placing in positions $i_1, \dots, i_k$ the values $\sigma_{i_2}, \dots, \sigma_{i_k}, \sigma_{i_1}$, respectively, and leaving all other entries of σ fixed. Let $\phi^*(\sigma)$ be obtained by applying ϕ repeatedly until no $k\dots 1$ -patterns remain. It is shown that ϕ^* induces a bijection from $S_n(k-1\dots 1k)$ onto $S_n(k\dots 1)$, and that from this bijection one can obtain an important ingredient of the proof, namely a bijection from $S_n(1\dots k)$ onto $S_n(k\dots 1)$.

In [3], Bousquet-Mélou and Steingrímsson prove the analogue of the Backelin-West-Xin Theorem in the context of involutions, and in so doing they must

prove that $\phi^*(\sigma^{-1}) = (\phi^*(\sigma))^{-1}$ for all permutations σ , so that the bijection from $S_n(1 \dots k)$ onto $S_n(k \dots 1)$ will also commute with inverses. The proof of the commutation result for ϕ^* is long and difficult, and Bousquet-Mélou and Steingrímsson ask for a reformulation of ϕ^* that will make this result obvious. In [6], Krattenthaler describes another bijection from $S_n(1 \dots k)$ into $S_n(k \dots 1)$, in terms of growth diagrams, and notes that this bijection clearly commutes with inverses. He asks what connection there is between this bijection and ϕ^* . In the present paper we answer both questions, by providing a reformulation of ϕ^* in terms of growth diagrams.

In proving their theorem, Backelin, West, and Xin find it necessary to work in the context of full rook placements on Ferrers boards, which includes permutations as a special case. For any Ferrers board F and any permutation τ , let $S_F(\tau)$ denote the set of all full rook placements on F that avoid τ . (The relevant definitions will be reviewed in Section 2.)

Backelin, West, and Xin prove that $|S_F(1 \dots k\rho)| = |S_F(k \dots 1\rho)|$, for all F and all permutations ρ of $\{k+1, \dots, k+\ell\}$, and in so doing they use an extension of ϕ^* to full rook placements. Bousquet-Mélou and Steingrímsson also use this extension, so they prove that ϕ^* commutes with inverses in this broader context. Accordingly, our reformulation of ϕ^* will be given in this context, or rather in the even broader context of arbitrary rook placements (not necessarily full), with the term “inverse” interpreted appropriately.

The outline of our paper is as follows. In Section 2 we review the needed background material on the Robinson-Schensted correspondence for partial permutations, Ferrers boards and rook placements, and growth diagrams. In Section 3 we give our reformulation of ϕ^* and the proof that it works, modulo a “Main Lemma”. In Section 4 we introduce the tool that will be used to prove this lemma: the “pivots” of a rook placement on a rectangular Ferrers board. The pivots are related to the “ L -corners” of [2], and are a generalization of the “ rcL -corners” of [8]. (We have chosen to use the term “pivots”, instead of “corners”, because of the prior use of the term “corners” in connection with the diagram of a permutation.) Section 5 contains the proof of the Main Lemma, and the concluding Section 6 indicates how the proof leads naturally to a notion of generalized Knuth transformations.

2. Review of the Needed Background

2.1. The Robinson-Schensted correspondence for partial permutations

A *partial permutation* π is a bijection between two sets I and J of positive integers. We represent π in two-line notation as

$$\pi = \begin{pmatrix} i_1 & i_2 & \cdots & i_m \\ j_1 & j_2 & \cdots & j_m \end{pmatrix}$$

where $i_1 < \dots < i_m$ and the j_t 's are distinct.

The Robinson-Schensted *insertion and recording tableaux* P and Q for π are obtained as follows. Start by placing j_1 in the top row of P and i_1 in the top row of Q . Assuming inductively that $i_1, \dots, i_{t-1}$ and $j_1, \dots, j_{t-1}$ have been placed, place j_t in P as follows. If j_t is larger than all elements already in the top row of P , place j_t at the right end of this row. If j_s is the leftmost element already in the top row that is larger than j_t , replace j_s by j_t and “bump” j_s to the second row. Place j_s at the right end of this row unless it is smaller than some element in this row, in which case let j_s bump the leftmost element larger than it to the third row, and continue in this way. Place i_t in the position in Q corresponding to the position in P that first became occupied when j_t was placed in P .

For example, if

$$\pi = \begin{pmatrix} 1 & 2 & 6 & 7 & 8 \\ 4 & 5 & 3 & 1 & 2 \end{pmatrix}$$

then

$$P_\pi = \begin{array}{|c|c|} \hline 1 & 2 \\ \hline 3 & 5 \\ \hline 4 & \\ \hline \end{array} \quad Q_\pi = \begin{array}{|c|c|} \hline 1 & 2 \\ \hline 6 & 8 \\ \hline 7 & \\ \hline \end{array}$$

A basic property of P and Q is that there exists a partition $\lambda = \{\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_t\}$ of m such that the top rows of P and Q each contain λ_1 entries, the second rows each contain λ_2 entries, and so on. The partition λ is called the *shape* of P and Q . In motivating our reformulation of ϕ^* it will be helpful to recall the theorem of Schensted [9] that states that λ_1 is the length of the longest increasing subsequence in π , and t is the length of the longest decreasing subsequence.

We will also use the theorem of Schützenberger [10] that states that the insertion and recording tableaux for the inverse bijection π^{-1} are Q and P , respectively.

2.2. Ferrers boards, rook placements, and ϕ^*

Consider an $n \times n$ array of squares, and identify the pair (i, j) with the square located in the i th column from the left and the j th row from the bottom. For any square (i, j) in the array, let $R(i, j)$ denote the rectangle consisting of all squares (k, ℓ) such that $k \leq i$ and $\ell \leq j$. A *Ferrers board* (in French notation) is any subset F of such an array with the property that for all $(i, j) \in F$ we have $R(i, j) \subseteq F$. So for some t and some $\lambda_1 \geq \dots \geq \lambda_t$, the Ferrers board consists of the first λ_j squares from the j th row of the array, $1 \leq j \leq t$. The *conjugate* of F is the Ferrers board $F' = \{(j, i) : (i, j) \in F\}$, so that F' is obtained by reflecting F across the *SW-NE* diagonal.

A *rook placement* on a Ferrers board F is a subset of F that contains at most one square from each row of F and at most one square from each column of F . We indicate the squares in the placement by putting markers (e.g., dots or X 's) in them. A rook placement is called *full* if it includes exactly one square from

each row and column of F . (So if there exists a full rook placement on F , then F has the same number of columns as rows.) From any rook placement P on F there results a partial permutation π such that square (i, j) is in P if and only if j is the value of the bijection π at input i . P is a full placement if and only if F has n rows and n columns for some n and π is a permutation of $\{1, \dots, n\}$. For any rook placement P on F , the *inverse* P' of P is the placement on the conjugate board F' obtained by reflecting F and all the markers for P across the SW - NE diagonal. The partial permutations resulting from P and P' are inverses of each other, if we regard them as bijections between sets. If they are permutations, they are inverses in the usual sense.

We say that a rook placement P *contains* a permutation $\tau \in S_r$ if and only if the resulting partial permutation π contains a subsequence $\pi_{i_1} \dots \pi_{i_r}$ order isomorphic to τ such that there is a rectangular subboard of F that contains all the squares (i_j, π_{i_j}) . In this case we refer to the sequence of squares (i_j, π_{i_j}) as an *occurrence* of τ in P . We say that P *avoids* τ if P does not contain τ .

It is clear how to extend the definitions of ϕ and ϕ^* to rook placements, by using only the occurrences of $k \dots 1$ in P , in the sense of the preceding paragraph.

2.3. Growth diagrams


Our reformulation of ϕ^* will be accomplished by modifying Fomin's ([4, 5], see also [6]) construction of the growth diagram of a rook placement P on a Ferrers board F .

Fomin's construction assigns partitions to the corners of all the squares in F , using the markers of P , in such a way that the partition assigned to any corner either equals the partition to its left or is obtained from it by adding 1 to one entry, and the partition assigned to any corner either equals the partition below it or is obtained from this partition by adding 1 to one entry. We start by assigning the empty partition $\emptyset$ to each corner on the left and bottom edges of F . We then assign partitions to the other corners inductively. Assuming that the northwest, southwest, and southeast corners of a square (i, j) have been assigned partitions NW, SW , and SE , we assign to the northeast corner the partition NE determined by the following rules.


1. If $NW \neq SE$ then let $NE = NW \cup SE$, the partition whose i th entry is the maximum of the i th entries of NW and SE . (Here we regard the absence of an entry as the presence of an entry 0.)
2. If $SW \neq NW = SE$ then NW is obtained from SW by adding 1 to the i th entry of SW , for some i . We obtain NE from NW by adding 1 to the $(i + 1)$ th entry.
3. If $SW = NW = SE$ then we let $NE = SW$ unless the square (i, j) contains a marker, in which case we obtain NE from SW by adding 1 to the first entry.

Lemma 1. ([6], Theorem 5.2.4 or [11], Theorem 7.13.5). For any square (i, j) , the partition assigned to the northeast corner of (i, j) is the shape of the Robinson-Schensted tableaux for the partial permutation resulting from the restriction of P to the rectangle $R(i, j)$.

Example. Consider the following (full) rook placement P on the indicated Ferrers board F .



The growth diagram is (denoting each partition by juxtaposing its entries)



Note that the partial permutation obtained by restricting P to $R(8, 5)$ is

$$\pi = \begin{pmatrix} 1 & 2 & 6 & 7 & 8 \\ 4 & 5 & 3 & 1 & 2 \end{pmatrix}$$

and the partition assigned to the northeast corner of square $(8, 5)$ is 221, which is, by the example given in subsection 2.1, the shape of the Robinson-Schensted tableaux for π .

In general, if we are given F and the partitions in the growth diagram for P that occur along the right/up border of F (i.e., the border of F minus the horizontal bottom edge and the vertical left edge), we can inductively reconstruct the rest of the growth diagram and the placement P by using the following rules to assign a partition to the southwest corner of square (i, j) , given the partitions assigned to its other three corners.

- A. If $NW \neq SE$ then let $SW = NW \cap SE$, the partition whose i th entry is the minimum of the i th entries of NW and SE .
- B. If $NW = NE = SE$ then let $SW = NW$.
- C. If $NE \neq NW = SE$ and NE differs from NW in the i th entry for some $i \geq 2$, then let SW be obtained from NW by subtracting 1 from the $(i - 1)$ th entry of NW . If NE differs from NW in the first entry, then let $SW = NW$ and, *in this circumstance only*, place a marker in square (i, j) .

Since the application of these rules recovers the growth diagram and the placement P , it follows that, in the growth diagram, square (i, j) has a marker in it (i.e., is in P) if and only if $NE \neq NW = SE$ and NE differs from NW in the first entry.

3. The Reformulation of ϕ^*

We now modify the growth diagram algorithm (GDA) of subsection 2.3 to get a new algorithm GDA_k for any $k \geq 2$. GDA_k retains rules (1) and (3) of GDA , but replaces rule (2) by the following variant.

- 2_k . Apply rule (2) with the proviso that if rule (2) produces a NE with k (nonzero) entries then delete the last entry and increase the first entry by 1.

The motivation for rule (2_k) comes from the theorem of Schensted mentioned in subsection 2.1. Keeping the number of entries in a partition λ less than k prevents decreasing subsequences of length k in partial permutations whose Robinson-Schensted tableaux have shape λ .

Definitions. For any rook placement P on a Ferrers board F , let $\text{seq}(P, F)$ (respectively, $\text{seq}_k(P, F)$) denote the sequence of partitions along the right/up border of F that results from the application of GDA (respectively, GDA_k) to P and F .

Main Theorem. Fix $k \geq 2$ in the definition of ϕ^* . Then for any rook placement P on a Ferrers board F ,

$$\text{seq}_k(P, F) = \text{seq}(\phi^*(P), F).$$

Corollary. For any rook placement P on a Ferrers board F ,

$$\phi^*(P') = (\phi^*(P))'.$$

Proof. This is essentially clear from the fact that the algorithms GDA and GDA_k commute with the operation of taking the inverse of a placement.

In a bit more detail, $\text{seq}_k(P', F')$ is the reverse of $\text{seq}_k(P, F)$, so, by the Main Theorem, $\text{seq}(\phi^*(P'), F')$ is the reverse of $\text{seq}(\phi^*(P), F)$, and this reverse is $\text{seq}((\phi^*(P))', F')$. By rules (A), (B), and (C) for the inverse algorithm for *GDA*, we conclude that $\phi^*(P') = (\phi^*(P))'$. $\square$

We will now give an example, to illustrate the Main Theorem and indicate the structure of its proof.

Example. Let P be the placement from the example in subsection 2.3, and fix $k = 3$ in the definition of ϕ^* . Performing GDA_3 on P yields

and performing *GDA* on $\phi^*(P)$ yields

$\emptyset$	1	2	• 3						
$\emptyset$	1	2	2	3	• 4				
$\emptyset$	1	2	2	• 3	3	31			
$\emptyset$	1	2	2	2	2	21	22	• 32	
$\emptyset$	1	• 2	2	2	2	21	22	22	
$\emptyset$	• 1	1	1	1	1	11	21	21	
$\emptyset$	$\emptyset$	$\emptyset$	$\emptyset$	$\emptyset$	$\emptyset$	1	• 2	2	
$\emptyset$	$\emptyset$	$\emptyset$	$\emptyset$	$\emptyset$	$\emptyset$	• 1	1	1	
$\emptyset$	$\emptyset$	$\emptyset$	$\emptyset$	$\emptyset$	$\emptyset$	$\emptyset$	$\emptyset$	$\emptyset$	

The partitions along the right/up border of F are the same in both cases, so $\text{seq}_3(P, F) = \text{seq}(\phi^*(P), F)$, although the partitions in the interiors of the diagrams are not always the same.

In this example, performing GDA_3 on $\phi(P)$ yields

$\emptyset$	1	2	• 3						
$\emptyset$	1	2	2	3	• 4				
$\emptyset$	1	2	2	• 3	3	31			
$\emptyset$	1	• 2	2	2	2	21	22	32	
$\emptyset$	1	1	1	1	1	11	• 21	22	
$\emptyset$	• 1	1	1	1	1	11	11	21	
$\emptyset$	$\emptyset$	$\emptyset$	$\emptyset$	$\emptyset$	$\emptyset$	1	1	• 2	
$\emptyset$	$\emptyset$	$\emptyset$	$\emptyset$	$\emptyset$	$\emptyset$	• 1	1	1	
$\emptyset$	$\emptyset$	$\emptyset$	$\emptyset$	$\emptyset$	$\emptyset$	$\emptyset$	$\emptyset$	$\emptyset$	

Notice that, while the results of performing GDA_3 on P and on $\phi(P)$ are not the same diagram, they do agree on the boundary of $R(7, 4)$, the smallest rectangular subboard of F that contains the 321-pattern on which ϕ acted and extends to the left and bottom edges of F . Because of the definition of GDA_3 , this is enough to make the two diagrams agree everywhere outside the rectangle. The idea of the proof of the Main Theorem will be to show that performing GDA_k on P and on $\phi(P)$ yields the same partitions on the boundary of the smallest rectangular subboard of F that contains the $k \dots 1$ -pattern on which ϕ acted and extends to the left and bottom edges of F .

Proof of the Main Theorem. We proceed by induction on the number of applications of ϕ required to compute $\phi^*(P)$. If no applications are required, the result is obvious, since $\phi^*(P) = P$ and performing GDA_k is the same as performing GDA .

Now suppose that m applications of ϕ are required to compute $\phi^*(P)$. Since computing $\phi^*(\phi(P))$ requires only $m-1$ applications of ϕ , we assume inductively that

$$\text{seq}_k(\phi(P), F) = \text{seq}(\phi^*(\phi(P)), F),$$

i.e., that

$$\text{seq}_k(\phi(P), F) = \text{seq}(\phi^*(P), F).$$

We want to show that $\text{seq}_k(\phi(P), F) = \text{seq}_k(P, F)$.

Let $R = R(a, b)$ be the smallest rectangular subboard of F that contains the $k \dots 1$ -pattern on which ϕ acted to produce $\phi(P)$ and extends to the left and bottom edges of F . Let P_R and $\phi(P)_R$ be the restrictions of P and $\phi(P)$ to $R(a, b)$. By the definition of GDA_k , all we need to show is that

$$\text{seq}_k(\phi(P)_R, R) = \text{seq}_k(P_R, R).$$

To show this, we will use the following two lemmas, which will be proved at the end of this section.

Lemma 2. The placement P_R on R contains no $k \dots 1$ -pattern that begins in a row below the top row of R or ends in a column to the left of the rightmost column of R .

Lemma 3. The placement $\phi(P)_R$ on R contains no $k \dots 1$ -pattern.

By Lemma 3,

$$\text{seq}_k(\phi(P)_R, R) = \text{seq}(\phi(P)_R, R).$$

By Lemma 2, $\text{seq}_k(P_R, R) = \text{seq}(P_R, R)$ except possibly at the northeast corner of square (a, b) .

Notation. Let c_{ne} , c_{nw} , and c_{se} denote the northeast, northwest, and southeast corners of square (a, b) .

To conclude the proof, it will suffice to prove the next lemma.

Main Lemma. We have $\text{seq}(P_R, R) = \text{seq}(\phi(P)_R, R)$ except possibly at c_{ne} , the northeast corner of R .

For once this lemma is established, we will have

$$\text{seq}_k(\phi(P)_R, R) = \text{seq}_k(P_R, R)$$

except possibly at c_{ne} . To see that the two also agree at c_{ne} , let, by the Main Lemma, λ be the common value of $\text{seq}(P_R, R)$ and $\text{seq}(\phi(P)_R, R)$ at c_{nw} . Since

$$\text{seq}_k(\phi(P)_R, R) = \text{seq}(\phi(P)_R, R),$$

the value of $\text{seq}_k(\phi(P)_R, R)$ at c_{ne} is obtained from λ by adding 1 to the first entry. (This follows from the last statement in subsection 2.3 and the fact that $\phi(P)_R$ has a marker in square (a, b) .) To see that $\text{seq}_k(P_R, R)$ has the same value at c_{ne} , let μ, ν be the values of $\text{seq}(P_R, R)$ at c_{se}, c_{ne} . Since, in the placement P_R on R , $R(a-1, b)$ and $R(a, b-1)$ contain no $k \dots 1$ -patterns (by Lemma 2) but $R(a, b)$ contains such a pattern, it must be that ν has k entries and each of λ, μ has $k-1$. Therefore, by the definition of GDA_k , the value of $\text{seq}_k(P_R, R)$ at c_{ne} is obtained by adding 1 to the first entry of λ .

The proof of the Main Theorem is now complete, except for the proofs of Lemmas 2 and 3 and the Main Lemma. The proof of the Main Lemma will occupy Section 5. We now turn to the proofs of Lemmas 2 and 3.

Notation. For a square $B = (i, j)$ in a Ferrers board F , we denote i and j by $\text{col}(B)$ and $\text{row}(B)$, respectively.

Proof of Lemma 2. Let $D = D_1 D_2 \dots D_k$ be the sequence of squares, from left to right, constituting the smallest occurrence of $k \dots 1$ in P . Let $S = S_1 S_2 \dots S_k$ be another occurrence of $k \dots 1$ in P . By the minimality of D , we cannot have $\text{row}(S_1) < \text{row}(D_1)$, so S cannot begin in a row below the top row of $R = R(a, b)$, and $S_1 = D_1$. Suppose for a contradiction that S ends in a column to the left of the rightmost column of R , so $\text{col}(S_k) < \text{col}(D_k)$.

If $\text{row}(S_k) > \text{row}(D_k)$ then $S_2 \dots S_k D_k$ contradicts the minimality of D . Clearly $\text{row}(S_k) \neq \text{row}(D_k)$, because S_k and D_k are in different columns. So $\text{row}(S_k) < \text{row}(D_k)$. Therefore $S_k \neq D_{k-1}$, so $\text{col}(S_k) \neq \text{col}(D_{k-1})$. If $\text{col}(S_k) > \text{col}(D_{k-1})$ then $D_1 \dots D_{k-1} S_k$ contradicts the minimality of D , since $\text{row}(S_k) < \text{row}(D_k) < \text{row}(D_{k-1})$. So $\text{col}(S_k) < \text{col}(D_{k-1})$.

Now, assuming inductively that $\text{col}(S_t) < \text{col}(D_{t-1})$ for some $t \geq 3$, we have $\text{col}(S_{t-1}) < \text{col}(D_{t-1})$. We will show by essentially the same argument as in the preceding paragraph that $\text{col}(S_{t-1}) < \text{col}(D_{t-2})$. First, if $\text{row}(S_{t-1}) > \text{row}(D_{t-1})$ then $S_2 \dots S_{t-1} D_{t-1} \dots D_k$ contradicts the minimality of D . So $\text{row}(S_{t-1}) < \text{row}(D_{t-1})$. If $\text{col}(S_{t-1}) > \text{col}(D_{t-2})$ then $D_1 \dots D_{t-2} S_{t-1} \dots S_k$ contradicts the minimality of D . Thus $\text{col}(S_{t-1}) < \text{col}(D_{t-2})$.

We conclude by induction that $\text{col}(S_2) < \text{col}(D_1) = \text{col}(S_1)$, a contradiction.

□

Proof of Lemma 3. Again let D be the smallest occurrence of $k \dots 1$ in P , and let $D_1^*, \dots, D_k^*$ be the squares by which $D_1, \dots, D_k$ are replaced in $\phi(P)_R$. Assume for a contradiction that $\phi(P)_R$ has a $k \dots 1$ -pattern, S , in R . Let c be the number of squares that $D^* = D_2^* \dots D_k^*$ has in common with S . (Note that D_1^* is the square in the northeast corner of R .) Let $d = k - c$ be the number of squares in S that are not in D^* . Observe that $c \geq 1$. If this were not the case then S would contain no square D_i^* and therefore it would also be a $k \dots 1$ -pattern in P_R . Then $\text{row}(S_1) < \text{row}(D_1^*) = \text{row}(D_1)$ would contradict the minimality of D .

Now let the c common squares, from left to right, be $D_{m_j}^*$ for $1 \leq j \leq c$. We group the other squares of S according to their rows. Let d_0 be the number of these other squares that are north of $D_{m_1}^*$. Let d_s be the number of the other squares that are south of $D_{m_s}^*$ and north of $D_{m_{s+1}}^*$, for $1 \leq s \leq c-1$. Finally let d_c be the number of the other squares that are south of $D_{m_c}^*$.

Note that all the squares counted by d_c must be to the right of $D_{m_c}^*$, so we must have $d_c \leq k - m_c$ because if there were $k - m_c + 1$ squares B_i counted by d_c then the sequence $D_1 \dots D_{m_{c-1}}$ followed by the B_i s would contradict the minimality of D . Next, we must have $d_s \leq m_{s+1} - m_s - 1$. To see this note that the squares counted by d_s must be west of the column containing $D_{m_{s+1}}^*$ and east of the column containing $D_{m_s}^*$. Therefore if there were $m_{s+1} - m_s$ squares B_i counted by d_s the sequence $D_1 D_2 \dots D_{m_{s-1}}$, followed by the B_i s and then $D_{m_{s+1}} \dots D_k$ would contradict the minimality of D . Finally, observe that $d_0 \leq m_1 - 2$. For if there were $m_1 - 1$ squares counted by d_0 then these B_i s followed by $D_{m_1} \dots D_k$ would contradict the minimality of D .

Now when $c > 1$ we have

$$\begin{aligned}
k &= c + d \\
&= c + d_0 + d_c + \sum_{s=1}^{c-1} d_s \\
&\leq c + (m_1 - 2) + (k - m_c) + \sum_{s=1}^{c-1} m_{s+1} - m_s - 1 \\
&\leq (c - 1) + (k - 1) + m_1 - m_c + m_c - m_1 - (c - 1) \\
&= k - 1.
\end{aligned}$$

When $c = 1$ we obtain the similar contradiction

$$k = 1 + d_0 + d_1 \leq 1 + (m_1 - 2) + (k - m_1) = k - 1.$$

Therefore $\phi(P)_R$ cannot contain a $k \dots 1$ -pattern in R , as claimed. $\square$

4. Pivots

In this section we introduce the *left* and *right pivots* of rook placement on a rectangular Ferrers board, and show how they relate to the Robinson-Schensted

correspondence and to ϕ^* . While the right pivots have nicer properties in connection with the RS correspondence, the left pivots have nicer properties with respect to ϕ^* .

Definitions. Let P be a placement on rectangular Ferrers board F . We define the set of *left pivots of P* (respectively, *right pivots of P*) to be the placement $\text{piv}_l P$ (respectively, $\text{piv}_r P$) defined by inductively placing markers, row by row, from bottom to top, as follows.

First, there is no pivot in the bottom row. Now consider row $r > 1$. If there is no element of P in row r then we do not place a pivot in row r . Now suppose $X \in P$ is in row r . If there is a column to the left (respectively, right) of X that contains an element of P below row r but does not contain a pivot then we place a pivot in row r and in the rightmost (respectively, leftmost) such column.

Notation. For a placement P on a rectangular Ferrers board F define $\text{rev}(P)$ to be the placement obtained by reflecting F and P along a vertical line.

Below is an example of left and right pivots of a placement.




Figure 1: On the left we have P and $\text{piv}_l(P)$. On the right we have $\text{rev}(P)$ and $\text{piv}_r(P)$. In both examples elements of P are denoted by $\times$ and the pivots are denoted by $\bullet$.

Looking at Figure 1 we see the following relationship between left and right pivots. Its proof is straightforward.

Lemma 4. For any placement P on a rectangular Ferrers board F we have

$$\text{rev}(\text{piv}_l P) = \text{piv}_r(\text{rev } P).$$

The utility of pivots is due to their connection with the Robinson-Schensted (RS) algorithm. We will consider applying the RS algorithm to a placement P . What we really mean is that we are implicitly applying it to the partial permutation corresponding to P . We will also write $\text{ins}(P)$ and $\text{rec}(P)$ for the

insertion and recording tableaux of P . Let us first compare the insertion and recording tableaux for P , $\text{piv}_r(P)$, and $\text{piv}_l(P)$ using the placement on the right hand side in Figure 1.

When we apply RS to P we get

$$\text{ins } P = \begin{array}{|c|c|c|} \hline 1 & 4 & 8 \\ \hline 2 & 5 & \\ \hline 7 & & \\ \hline 9 & & \\ \hline \end{array} \quad \text{rec } P = \begin{array}{|c|c|c|} \hline 1 & 3 & 9 \\ \hline 4 & 8 & \\ \hline 6 & & \\ \hline 7 & & \\ \hline \end{array}.$$

Now on the one hand RS applied to $\text{piv}_r P$ gives us

$$\text{ins}(\text{piv}_r P) = \begin{array}{|c|c|} \hline 2 & 5 \\ \hline 7 & \\ \hline 9 & \\ \hline \end{array} \quad \text{rec}(\text{piv}_r P) = \begin{array}{|c|c|} \hline 4 & 8 \\ \hline 6 & \\ \hline 7 & \\ \hline \end{array},$$

while RS applied to $\text{piv}_l P$ gives

$$\text{ins}(\text{piv}_l P) = \begin{array}{|c|c|} \hline 4 & 8 \\ \hline 5 & \\ \hline \end{array} \quad \text{rec}(\text{piv}_l P) = \begin{array}{|c|c|} \hline 1 & 8 \\ \hline 7 & \\ \hline \end{array}.$$

We see here that the RS algorithm applied to $\text{piv}_r P$ gives the same insertion and recording tableaux, minus the top row, as the RS algorithm applied to P . And the insertion tableau of $\text{piv}_l P$ is equal to $\text{ins } P$, minus its left column. This is the content of the next two theorems.

Notation. If Y is a standard Young tableau then we will denote by Y^- the tableau consisting of all but the top row of Y and ^-Y the tableau consisting of all but the left column of Y .

Theorem 1. Let P be a placement on a rectangular Ferrers board. Then we have

$$\text{ins}(\text{piv}_r P) = (\text{ins } P)^- \quad \text{and} \quad \text{rec}(\text{piv}_r P) = (\text{rec } P)^-$$

Proof. Let

$$\pi = \begin{pmatrix} i_1 & i_2 & \cdots & i_n \\ s_1 & s_2 & \cdots & s_n \end{pmatrix},$$

with $i_1 < i_2 < \dots$, be the partial permutation corresponding to P . Let

$$\begin{pmatrix} p_1 & p_2 & \cdots & p_m \\ v_1 & v_2 & \cdots & v_m \end{pmatrix},$$

with $p_1 < p_2 < \dots$, be the partial permutation corresponding to $\text{piv}_r(P)$.

If we consider the RS algorithm applied to π it will suffice to show that inserting $\pi(p_j)$ into the insertion tableau causes v_j to be bumped from the first row and that the v_j are the only numbers bumped from the first row. To show this we use induction on m , the number of pivots. If $m = 0$ then π must have

no 21-pattern and hence $s_1 < s_2 < \dots$. Hence $(\text{ins } \pi)^- = \emptyset$ and $(\text{rec } \pi)^- = \emptyset$. Now assume π is such that $m > 0$. Define k so that $i_k = p_m$ and consider $\pi|_{k-1}$, the restriction of π to its first $k-1$ entries. Then we have

$$\text{piv}_r(\pi|_{k-1}) = \begin{pmatrix} p_1 & p_2 & \cdots & p_{m-1} \\ v_1 & v_2 & \cdots & v_{m-1} \end{pmatrix}.$$

By induction the RS algorithm applied to $\pi|_{k-1}$ bumps v_j on the p_j th move for $1 \leq j \leq m-1$ and bumps nothing else. So the top row of the insertion tableau is just

$$S = \{s_j \mid 1 \leq j \leq k-1\} \setminus \{v_j \mid 1 \leq j \leq m-1\}.$$

Since applying the RS algorithm to $\pi|_{k-1}$ is the same as partially computing the RS algorithm on π up through the index i_{k-1} of π we may assume this is where we are in the algorithm. We now claim that inserting the next value of π , s_k , into the insertion tableau bumps v_m from the first row of the insertion tableau. Showing this is equivalent to showing that $v_m \in S$ and that v_m is the smallest element in S that is greater than s_k . Clearly $v_m \in S$ as all the v_j are distinct. Now if $s \in \{s_j \mid 1 \leq j \leq k-1\}$ and $s_k < s < v_m$ then ss_k is a 21-pattern. But this implies that for some $i < m$, $s = v_i$. Hence $s \notin S$ and v_m is indeed bumped by the insertion of s_k . Therefore after the k th move the first row of the insertion tableau is

$$S' = \{s_j \mid 1 \leq j \leq k\} \setminus \{v_j \mid 1 \leq j \leq m\}.$$

Now let $k < j \leq n$. To complete the proof we need only show that the RS algorithm appends s_j to the top row of the insertion tableau. This is equivalent to showing that $s_{k+1} < s_{k+2} < \dots$ and that s_{k+1} is greater than all the elements in S' . If the s_j are not increasing from $j = k+1$ onward then they must contain a 21-pattern and hence there exists a pivot to the right of position p_m , contradicting our choice of p_m . If s_{k+1} is not greater than all the elements in S' then for some $s' \in S'$, $s's_{k+1}$ a 21-pattern. This means that either $s' = v_i$, for some i , or $i_{k+1} = p_j$, for some j . But this contradicts either our definition of S' or our definition of p_m . $\square$

Remark 1. We thank Sergi Elizalde for suggesting to us that there might be a connection between our right pivots and Viennot's geometric construction. It turns out that the right pivots coincide with Viennot's "northeast corners". This follows from the fact that Viennot establishes the analogue of Theorem 1 for the northeast corners.

To state our theorem relating left-pivots and the RS algorithm we first need the following.

Notation. If Y is any standard Young tableau then denote by Y^{tr} the transposed tableau, i.e., the tableau obtained by reflecting Y across the NW-SE diagonal.

Likewise, if P is any placement on a rectangular Ferrers board F then denote by F^{tr} and P^{tr} the resulting board and placement obtained by reflecting F and P across the NW-SE diagonal.

Remark 2. Recall the well known fact that for any placement P we have

$$\text{ins}(\text{rev } P) = (\text{ins } P)^{tr}.$$

For a proof of this result see [7], page 97.

Theorem 2. Let P be a placement on a rectangular Ferrers board F . Then

$$\text{ins}(\text{piv}_l P) = {}^-(\text{ins } P).$$

Proof. By Lemma 4 we have

$$\text{rev}(\text{piv}_l P) = \text{piv}_r(\text{rev } P).$$

Taking the insertion tableau of both sides and applying Theorem 1 to the right hand side we get

$$\text{ins}(\text{rev}(\text{piv}_l P)) = [\text{ins}(\text{rev } P)]^-.$$

Then by Remark 2 we have

$$(\text{ins}(\text{piv}_l P))^{tr} = [(\text{ins } P)^{tr}]^-.$$

By then transposing both sides we have

$$\text{ins}(\text{piv}_l P) = ([(\text{ins } P)^{tr}]^-)^{tr} = {}^-(\text{ins } P).$$

□

Definitions. Let P be a placement on a rectangular Ferrers board. For any $X \in P$, if X is in the same column as some $V \in \text{piv}_l P$ then define $\rho(P, X) = \text{row}(V)$ else define $\rho(P, X) = \infty$.

Likewise, if X is in the same row as some $V \in \text{piv}_l P$ then define $\kappa(P, X) = \text{col}(V)$ else define $\kappa(P, X) = 0$.

If the placement is understood we will just write $\rho(X)$ or $\kappa(X)$.

Next let us establish some symmetry intrinsic to left-pivots

Lemma 5. If P is a placement on a rectangular Ferrers board F with n columns, then we have

$$(\text{piv}_l P)^{tr} = \text{piv}_l(P^{tr})$$

Proof. Observe that this is equivalent to showing that the set $\text{piv}_l(P)$ may be constructed by the following column construction.

Column Construction

First, there is no pivot in the rightmost column. Now consider column $c < n$. If there is no element of P in column c then we do not place a pivot in column c . If $X \in P$ is in column c then if there is a row above X that contains no pivot to the right of column c but does contain an element of P to the right of column c then there is a pivot in column c and in the lowest such such row.

For reference we will call these pivots *column pivots*. To show that the column and row constructions agree assume inductively that

$$\text{column pivots}(P) = \text{piv}_l(P), \text{ for all columns to the right of } c. \quad (1)$$

Assume for a contradiction that the column pivots and left pivots disagree in column c . Note that if there is no column pivot in column c then by (1) we cannot have a left pivot either. Now assume we have a column pivot in (c, r) , for some r . So there must exist $X, Y \in P$ with $\text{col}(X) = c$, $\text{row}(Y) = r$ and XY a 12-pattern. If there is no left pivot in (c, r) then it follows from (1) that $\kappa(Y) < c$. But this can only occur if there exists some $Z \in P$ with $\text{row}(Z) < r$ and $\kappa(Z) = c$. But then (1) and the column construction imply that the column pivot in column c would be in some row $< r$ and not in row r as assumed. $\square$

The following definition and lemmas will be useful in Sections 5 and 6.

Convention. Since we will only be working with left pivots for the remainder of this paper, the term pivot will mean left pivot from now on.

Definition. We say an increasing subsequence I of P is a *pivot-path*, if for all $i < |I|$, $\rho(I_i) = \text{row}(I_{i+1})$, i.e., each consecutive pair creates a pivot.




Figure 2: An example of pivot-paths where the placement is denoted by $\times$ and the pivots are denoted by $\bullet$. The consecutive elements of each pivot path are connected by red lines.

Lemma 6. If $X, Y \in P$ are such that XY is a 12-pattern then we cannot have $\text{row}(Y) < \rho(X)$ and $\kappa(Y) < \text{col}(X)$.

Proof. If both of these held, then according to the construction of pivots we would have $\text{col}(X) \leq \kappa(Y)$. $\square$

Lemma 7. Let $X, Y, Z \in P$ be such that XYZ is a 132-pattern. If there is a pivot-path I with first element X and last element Y then there exists a pivot-path J with last element Z such that XJ_1 is a 21-pattern.

Proof. Let K be the longest pivot-path with last element Z . Note that $K \cap I = \emptyset$, for if not then $K \cup I$ is a pivot-path but this is prohibited as XYZ is a 132-pattern. Observe that I and K cannot “cross”, i.e., for all 12-patterns $I_i K_j$, we cannot have $\text{row}(K_j) < \rho(I_i)$ and $\kappa(K_j) < \text{col}(I_i)$ by Lemma 6. Therefore, if XK_1 is a 12-pattern then for some i , $I_i K_1$ is a 12-pattern with $\text{row}(K_1) < \rho(I_i)$. But by the maximality of the length of K we must have $\kappa(K_1) = 0 < \text{col}(I_i)$. But this cannot happen by Lemma 6. Therefore, if K_1 is to the right of X we can take $J = K$. If K_1 is to the left of X , define K_l to be the first element of K that is to the right of X . Observe that K_l must be below X since J and I do not cross. Take $J = K_l K_{l+1} \dots Z$. $\square$

The statement of the next lemma is similar to Lemma 7. Its proof is also similar to that of Lemma 7, but simpler.

Lemma 8. Let $X, Y \in P$ such that XY is a 12-pattern. If $\text{row}(Y) < \rho(X)$ then there exists a pivot-path J , with last element Y , such that XJ_1 is a 21-pattern.

Lemma 9. Let S be a decreasing sequence of length k in P . Assume there is some $X \in P$ such that XS_1 is a 12-pattern. If $\rho(X) > \text{row}(S_1)$ there exists a decreasing sequence D of length $k + 1$ with $D_1 = X$, and if X and S_1 are the first and last elements of a pivot-path then there exists a decreasing sequence D of length k with $D_1 = X$.

Proof. First consider the case when X and S_1 are connected by a pivot-path. We will proceed by induction on k , the length of S . If $k = 1$ then we may take $D = X$. Now consider $k > 1$. If $\text{row}(X) > \text{row}(S_2)$ then we may take $D = XS_2 \dots S_k$. If on the other hand $\text{row}(X) < \text{row}(S_2)$ then Lemma 7 implies that there exists a pivot-path J with last element S_2 and XJ_1 a 21-pattern. Now by the induction hypothesis applied to J_1 and the shorter sequence $S_2 S_3 \dots S_k$ we have a decreasing sequence E of length $k - 1$ with $E_1 = J_1$. But then the decreasing sequence $D = XE_1 E_2 \dots E_{k-1}$ is what we want.

For the case when $\rho(X) > \text{row}(S_1)$ first apply Lemma 8 to the 12-pattern XS_1 to obtain a pivot-path J such that XJ_1 forms a 21-pattern and J 's last element is S_1 . Now apply the first case to J_1 and the sequence S to obtain a decreasing sequence E of length k with $E_1 = J_1$. Then we may take $D = XE_1 E_2 \dots E_k$ which is of length $k + 1$. $\square$

Note that the previous lemma and Lemma 5 directly imply the following result.

Lemma 10. Let S be a decreasing sequence of length k in P . Assume there is some $X \in P$ such that $S_1 X$ is a 12-pattern. If $\kappa(X) < \text{col}(S_1)$ then there exists a decreasing sequence D of length $k + 1$ with $D_1 = X$.

5. The Proof of the Main Lemma

The purpose of this section is to prove the Main Lemma which is the crucial piece needed to show that our growth diagram construction corresponds to the map ϕ^* . Note that up through Lemma 11, F will always denote a rectangular Ferrers board and P a rook placement on F .

We begin with some notation and definitions.

Notation. Define $F|_{a,b}$ to be the Ferrers board consisting of all columns of F between and including columns a and b , and define $P|_{a,b}$ to be the restriction of P to $F|_{a,b}$.

Definitions. Fix a subplacement S , let $k = |S|$, and order S such that $S = S_1 \dots S_k$ where $\text{col}(S_i) < \text{col}(S_{i+1})$. Further fix a column c and assume that P has no element in column c .

We first define a *left-shift*. Assume S lies entirely to the right of column c . Place markers in all the squares of P , and then shift each marker in square S_i for $1 < i \leq k$ horizontally left to column $\text{col}(S_{i-1})$ and shift the marker in S_1 horizontally left to column c . Define $P(c \leftarrow S)$ to be the squares that now contain markers. Likewise, define $c \leftarrow S$ to be $P(S \leftarrow c) \setminus P$, the placement obtained from S by shifting it left.

Analogously, we define a *right-shift*. Assume S lies entirely to the left of column c . Place markers in all the squares of P , and then shift each marker in square S_i for $1 \leq i < k$ horizontally right to column $\text{col}(S_{i+1})$ and shift the marker in S_k horizontally right to column c . Define $P(S \rightarrow c)$ to be the squares that now contain markers. Likewise, define $S \rightarrow c$ to be $P(S \rightarrow c) \setminus P$, the placement obtained from S by shifting it right.

Definitions. Let P be a placement on a rectangular Ferrers board F . Let $a < b$ be columns of F .

Let k be the length of the longest decreasing sequence in $P|_{a,b}$. Define $\mathbf{d}_{a,b}(P)$ (respectively, $\mathbf{D}_{a,b}(P)$) to be the smallest (respectively, largest) decreasing sequence, under the lexicographical ordering, in $P|_{a,b}$ of length k .

Likewise, let m be the length of the longest increasing sequence in $P|_{a,b}$. Define $\mathbf{i}_{a,b}(P)$ (respectively, $\mathbf{I}_{a,b}(P)$) to be the smallest (respectively, largest) increasing sequence, under the lexicographical ordering, in $P|_{a,b}$ of length m .

We are now ready to state and prove Lemma 11. Although Lemma 11 is the key driver behind the Main Lemma, its statement is more general than what is needed to prove the Main Lemma. The reason for this is that the precise statement of Lemma 11 is exactly what is needed in Section 6.

Lemma 11. Let P be a placement on a rectangular Ferrers board F and assume P has no marker in column a . Let $S = \mathbf{d}_{a,b}(P)$ and $P^* = P(a \leftarrow S)$. Then we have

$$\text{ins}(P) = \text{ins}(P^*).$$

Proof. First let $S^* = a \leftarrow S$, i.e., the shifted sequence, and $k = |S|$. Define b' to be the column containing S_k and consider the truncated board and placements $G = F|_{1,b'}$, $Q = P|_{1,b'}$ and $Q^* = P^*|_{1,b'}$. Now in order to show

$$\text{ins}(P) = \text{ins}(P^*),$$

it will suffice to show

$$\text{ins}(Q) = \text{ins}(Q^*). \quad (2)$$

To see this consider the GDA applied to P and P^* on the full board F . By [11], Theorem 7.13.5, knowing (2) is the same as knowing that the partitions along the line $x = b'$ in the GDA of P and in the GDA of P^* are identical. But the placements P and P^* are identical east of the line $x = b'$. Therefore if (2) holds we know that the partitions along the right border of F in the GDA of P and in the GDA of P^* are also identical. By [11], Theorem 7.13.5, this implies that

$$\text{ins}(P) = \text{ins}(P^*).$$

In order to show (2) it is sufficient to prove

$$\text{piv}_l Q = \text{piv}_l Q^*. \quad (3)$$

To see why note that (3) along with Theorem 2 implies that

$$^-[\text{ins}(Q)] = ^-[\text{ins}(Q^*)]$$

But this forces $\text{ins}(Q) = \text{ins}(Q^*)$ as needed.

To establish (3) we will proceed inductively by assuming that the pivots (if any) below some row r are unchanged and then showing that the pivot (if any) on row r is unchanged. To do so we consider two cases.

Case 1: Row r contains an element S_i of S .

Let R be the rectangular region of F containing all squares west of col S_i and south of row S_i . By the nature of a left-shift, $Q|_R = Q^*|_R$. Further, by our induction hypothesis $\text{piv}_l(Q|_R) = \text{piv}_l(Q^*|_R)$. It then follows that the only way the pivot (if any) on row r could change is if $\text{piv}_l(Q)$ contains a pivot in row r between columns a' and $\text{col}(S_i)$, where $a' = a$, if $i = 1$, and $a' = \text{col}(S_{i-1})$, if $i > 1$.

To show this is impossible assume, for a contradiction, that $\text{piv}_l(Q)$ does contain a pivot in row r between columns a' and $\text{col}(S_i)$. Then there must exist some $X \in Q$ that is directly below this pivot. Since XS_i is a pivot-path, Lemma

9, applied to X and $S_i S_{i+1} \dots$, implies the existence of a decreasing sequence D of length $k - i + 1$ with first element X . But this is a contradiction since the decreasing sequence $S_1 \dots S_{i-1} D$ is smaller than S .

Case 2: Row r contains $X \in Q \setminus S$.

By the maximality of the length of S , $S_{i-1} X S_i$ cannot be a 321-pattern for any i . Therefore it will suffice to assume that $S_i X$ is a 12-pattern for some i and that i is chosen as small as possible. Then $\text{col}(S_i) \leq \kappa(X, Q)$. If not then Lemma 10 would give rise to sequence of length $k+1$ contradicting the maximality of the length of S . Therefore we must have an element $Y \in Q$ with $\text{col}(S_i) \leq \text{col}(Y) = \kappa(X, Q)$. Since $\text{col}(S_i) \leq \text{col}(Y) < b'$ then column $\text{col}(Y)$ must contain an element of Q^* below row r . (Note that for the case where $Y \in S$, Y cannot be the last element of S .) This plus the induction hypothesis implies that $\text{col}(Y) \leq \kappa(X, Q^*)$. Assume now, for a contradiction, that $\text{col}(Y) < \kappa(X, Q^*)$ and let $Z \in Q^*$ be such that $\text{col}(Y) < \text{col}(Z) = \kappa(X, Q^*)$. Now, by similar reasoning, $\text{col}(Z)$ must contain an element of Q below row r . But this plus the induction hypothesis implies that $\kappa(X, Q) = \text{col}(Y) < \text{col}(Z) \leq \kappa(X, Q)$, an obvious contradiction. $\square$

To prove the Main Lemma it suffices, by [11], Theorem 7.13.5, to prove the following.

Lemma 12. Let P be a placement on a not necessarily rectangular Ferrers board. Let S be the smallest $k \dots 1$ -pattern in P and let $b = \text{row}(S_1)$ and $a = \text{col}(S_k)$. If $R_1 = R(a, b-1)$ then

$$\text{ins}(P|_{R_1}) = \text{ins}(\phi(P)|_{R_1}). \quad (4)$$

Likewise, if $R_2 = R(a-1, b)$ then

$$\text{rec}(P|_{R_2}) = \text{rec}(\phi(P)|_{R_2}). \quad (5)$$

Proof. Let $Q = P|_{R_1}$ and $T = \phi(P)|_{R_1}$. Observe that $S_2, \dots, S_k = \mathbf{d}_{\text{col}(S_1), a}(Q)$. Now Lemma 11 implies (4).

For (5) let $R = R(a, b)$ and $Q = P|_R$ and $T = \phi(P)|_R$. Observe that $T' = \phi(Q')$, where Q' and T' are the inverses of Q and T in the sense of Section 2. This follows from the fact that S is the decreasing sequence of maximal length in R , which implies that S is also the position smallest sequence of length k in R (i.e., S is the smallest if we order $k \dots 1$ -patterns lexicographically according to the positions of the entries, rather than the values of the entries), so that S' is the value smallest decreasing sequence of length k in R' .




Figure 3: On the left we have an example of Q and T . On the right we have Q' and T' . In both pictures the $\phi(S)$ is marked by the red $\times$'s.

Applying (4), we know that

$$\text{ins}(Q'|_{R'_2}) = \text{ins}(T'|_{R'_2}).$$

So by the theorem of Schützenberger mentioned in Subsection 2.1 we have

$$\text{rec}(Q|_{R_2}) = \text{rec}(T|_{R_2}).$$

This completes the proof. $\square$

6. Generalized Knuth Transformations

Recall (see, for example, [11], page 414) that a *Knuth transformation* of a permutation σ interchanges two adjacent elements x and z of σ provided that there exists a third element y , which is adjacent to either x or z , such that $x < y < z$. Two permutations σ and ρ are called *Knuth-equivalent*, written $\sigma \sim_K \rho$, if ρ may be obtained from σ by a sequence of Knuth transformations.

It is a well known result that two permutations are Knuth-equivalent if and only if they share the same insertion tableau. For a proof of this result see, for example, [11], page 414.

Fix a placement P on a rectangular Ferrers board F . Further fix columns $a < b$ in F . Let $\mathbf{d} = \mathbf{d}_{a,b}(P)$, $\mathbf{D} = \mathbf{D}_{a,b}(P)$, $\mathbf{i} = \mathbf{i}_{a,b}(P)$, and $\mathbf{I} = \mathbf{I}_{a,b}(P)$.

Definition. Transforming the placement P into either of

$$P(a \leftarrow \mathbf{d}) \quad \text{or} \quad P(a \leftarrow \mathbf{I})$$

if P has no placement in column a , or to either of

$$P(\mathbf{D} \rightarrow b) \quad \text{or} \quad P(\mathbf{i} \rightarrow b)$$

if P has no placement in column b , is called a *generalized Knuth transformation*.

The next theorem, which is the main point of this section, shows that any two placements that differ by these generalized Knuth transformations are actually Knuth-equivalent.

Theorem 3. If two placements P and Q differ by generalized Knuth transformations then $\text{ins}(P) = \text{ins}(Q)$.

Before launching into the proof of this theorem let us pause to see how these four new transformations are generalizations of the standard Knuth transformations. First consider the Knuth transformation

$$\underbrace{5 \ 3 \ 4 \ 7 \ 6 \ 2 \ 8 \ 1}_{\sigma} \mapsto \underbrace{5 \ 3 \ 4 \ 7 \ 6 \ 2 \ 1 \ 8}_{\rho}.$$

Note that this is of the form $yzx \mapsto yxz$. Now consider P_{σ} to be the placement corresponding to σ on a 9×8 rectangular board where we have an empty column between the ‘6’ and the ‘2’. Observe that $\mathbf{d}_{6,9}(P_{\sigma})$ corresponds to the subsequence 21 in σ and that the partial permutation associated to

$$P_{\sigma}(6 \leftarrow \mathbf{d}_{6,9}(P_{\sigma}))$$

is order-isomorphic to ρ . It follows that any Knuth transformation of the form $yzx \mapsto yxz$ is generalized by the transformation $P \mapsto P(a \leftarrow \mathbf{d})$. Similar reasoning demonstrates how the remaining 3 standard Knuth transformations correspond by the remaining 3 generalized Knuth transformations. For completeness we give the correspondence in the table below.

Generalized	Standard
$P \mapsto P(a \leftarrow \mathbf{d})$	$\cdots yzx \cdots \mapsto \cdots yxz \cdots$
$P \mapsto P(\mathbf{D} \rightarrow b)$	$\cdots yxz \cdots \mapsto \cdots yzx \cdots$
$P \mapsto P(\mathbf{i} \rightarrow b)$	$\cdots xzy \cdots \mapsto \cdots zxy \cdots$
$P \mapsto P(a \leftarrow \mathbf{I})$	$\cdots zxy \cdots \mapsto \cdots xzy \cdots$

Observe that the standard Knuth transformations are clearly reversible. For example $yzx \mapsto yxz$ and $yxz \mapsto yzx$ are obviously inverses. This nice property of the standard transformations extends to the generalized transformations. We will see below in Lemma 15 that transformations of the forms $P(a \leftarrow \mathbf{d})$ and $P(\mathbf{D} \rightarrow b)$ are inverses and it follows by considering the reverse of a placement that transformations of the forms $P(\mathbf{i} \rightarrow b)$ and $P(a \leftarrow \mathbf{I})$ are inverses.

It remains to prove Theorem 3. This will occupy the remainder of this section. We start with a simple definition.

Definition. If S is a decreasing sequence in P then we say that a subplacement $A \subset P$ is *above* S if for every $C \in A$ there exists some index i such that $S_i C$ is a 12-pattern. Likewise, we say that $B \subset P$ is *below* S if for every $C \in B$ there exists some i such that $C S_i$ is a 12-pattern.

Lemma 13. Let L be a longest decreasing sequence in $P|_{a,b}$, of length k . If $L \neq \mathbf{D} = \mathbf{D}_{a,b}(P)$ then $P(L \rightarrow b)|_{a,b}$ contains a decreasing sequence of length $k + 1$. If $L \neq \mathbf{d}$ then $P(a \leftarrow L)|_{a,b}$ contains a decreasing sequence of length $k + 1$.

Proof. We prove only the first assertion. The second is analogous.

As $L \neq \mathbf{D}$ we must have some sequence D in $F|_{a,b}$ with $|D| = k$ that is value-larger than L . For D to be value-larger than L we must have some $0 \leq l$ and some $l + 1 \leq m$ such that

$$D = L_1 \dots L_l D_{l+1} \dots D_m D_{m+1} \dots D_k$$

where $D_{l+1} \dots D_m$ is above L and D_{m+1} is not above L , or D_m is the last element of D . Now define n to be the index such that D_m is between the columns containing L_n and L_{n+1} , or $n = k$ if D_m is the last element of D .

Clearly $D_{m+1} \neq L_n$ and as D_{m+1} is not above L then $L_n D_{m+1}$ must be a 21-pattern. Therefore

$$L_1 \dots L_n D_{m+1} \dots D_k$$

is a decreasing sequence in $P|_{a,b}$. Since no decreasing sequence in $P|_{a,b}$ has length greater than k then we must have $n + k - m \leq k$, i.e., $n \leq m$. Further, observe that

$$D_1 \dots D_m L_{n+1} \dots L_k$$

is also a decreasing sequence in $P|_{a,b}$. By similar logic we must have $m \leq n$. Therefore $m = n$.

Now consider the right-shift and let $L^* = L \rightarrow b$. Finally, observe that

$$L_1^* \dots L_l^* D_{l+1} \dots D_m L_n^* \dots L_k^*$$

is a decreasing sequence in $P(L \rightarrow b)|_{a,b}$. Since $m = n$ its length is $k + 1$ as claimed. $\square$

Lemma 14. We have the following relationships for decreasing sequences:

$$\begin{aligned} a \leftarrow \mathbf{d} & \text{ is longest in } P(a \leftarrow \mathbf{d})|_{a,b} \\ \mathbf{D} \rightarrow b & \text{ is longest in } P(\mathbf{D} \rightarrow b)|_{a,b} \end{aligned}$$

Proof. First let $Q = P|_{a,b}$ and define $n = b - a + 1$, the number of columns in Q . Now define

$$Q_1 = Q(1 \leftarrow \mathbf{d}_{1,n}(Q)) \qquad Q_2 = Q(\mathbf{D}_{1,n}(Q) \rightarrow n).$$

It will suffice, by the result of Schensted mentioned in Subsection 2.1, to prove that

$$\text{shape}(Q_i) = \text{shape}(Q).$$

For the first assertion, $\text{shape}(Q_1) = \text{shape}(Q)$ follows directly from Lemma 11.

For the second assertion let Q^* and Q_2^* be the complements of the reverses of Q and Q_2 . (By the complement of Q we mean the placement obtained by flipping Q and $F|_{a,b}$ about a horizontal line.) Note that

$$Q_2^* = Q^*(1 \leftarrow \mathbf{d}_{1,n}(Q^*)).$$

From the first assertion we know that $\text{shape}(Q_2^*) = \text{shape}(Q^*)$. Therefore $\text{shape}(Q_2) = \text{shape}(Q)$. $\square$

The next lemma will enable us to reverse generalized Knuth transformations. We state the results explicitly only for decreasing sequences. As remarked previously the results for increasing sequences follow immediately.

Lemma 15. Let $k = |\mathbf{d}|$ and assume that $\mathbf{d}_k$ is in column b . Then we have

$$a \leftarrow \mathbf{d} = \mathbf{D}_{a,b}(P(a \leftarrow \mathbf{d})).$$

Likewise, if $\mathbf{D}_1$ is in column a , then we have

$$\mathbf{D} \rightarrow b = \mathbf{d}_{a,b}(P(\mathbf{D} \rightarrow b)).$$

Proof. For the first claim set $P^* = P(a \leftarrow \mathbf{d})$. Assume for a contradiction that $a \leftarrow \mathbf{d} \neq \mathbf{D}_{a,b}(P^*)$. By Lemma 14 we know that $a \leftarrow \mathbf{d}$ is a longest sequence in $P(a \leftarrow \mathbf{d})|_{a,b}$. Now Lemma 13 implies that $P^*((a \leftarrow \mathbf{d}) \rightarrow b)|_{a,b} = P|_{a,b}$ has a decreasing subsequence of length $k + 1$. But this is impossible since $|\mathbf{d}| = k$.

A similar argument may be used to obtain the second claim. The details are straightforward. $\square$

Proof of Theorem 3. It will suffice to prove that the insertion tableau of a placement is invariant under any of the four generalized Knuth transformations. We give the arguments for decreasing sequences only.

Case 1: $P(a \leftarrow \mathbf{d})$.

This is clear by 11.

Case 2: $P(\mathbf{D} \rightarrow b)$.

First define a' to be the column containing the first element of $\mathbf{D}$. Lemma 15 implies that $\mathbf{D} \rightarrow b = \mathbf{d}_{a',b}[P(\mathbf{D} \rightarrow b)]$. Letting $P^* = P(\mathbf{D} \rightarrow b)$, Case 1 gives

$$\text{ins } P^* = \text{ins}[P^*(a' \leftarrow (\mathbf{D} \rightarrow b))] = \text{ins } P.$$

$\square$

Acknowledgment. The first author would like to thank his thesis advisor, Sergi Elizalde, for suggesting the problem of reformulating ϕ^* .

References

1. J. Backelin, J. West, G. Xin, Wilf-equivalence for singleton classes, *Adv. Appl. Math.* **38** (2007), 133–148.
2. J. Bloom and D. Saracino, Another look at pattern-avoiding permutations, *Adv. Appl. Math.* **45** (2010), 395–409.
3. M. Bousquet-Mélou, E. Steingrímsson, Decreasing subsequences in permutations and Wilf equivalence for involutions, *J. Algebraic Combin.* **22** (2005), 383–409.
4. S. Fomin, Generalized Robinson-Schensted-Knuth correspondence, *Zapiski Nauchn. Sem. LOMI* **155** (1986), 156–175.
5. S. Fomin, Schensted algorithms for dual graded graphs, *J. Alg. Comb.* **4** (1995), 5–45.
6. C. Krattenthaler, Growth diagrams, and increasing and decreasing chains in fillings of Ferrers shapes, *Adv. Appl. Math.* **37** (2006), 404–431.
7. B. Sagan, The Symmetric Group, second edition, Springer-Verlag, New York 2001.
8. D. Saracino, On two bijections from $S_n(321)$ to $S_n(132)$, *Ars. Comb.* **101** (2011), 65–74.
9. C. Schensted, Longest increasing and decreasing subsequences, *Canad. J. Math* **13** (1961), 179–191.
10. M. Schützenberger, Quelques remarques sur une construction de Schensted, *Math. Scand.* **12** (1963), 117–128.
11. R. Stanley, Enumerative Combinatorics, Vol. II, Cambridge Univ. Press, Cambridge, 1999.